



UNIVERSITÉ
DE GENÈVE



ÉCOLE POLYTECHNIQUE
FÉDÉRALE DE LAUSANNE

An architect gene is involved in the assimilation of breast milk

The presence of a mutated Hoxd3 gene could explain certain types of intestinal insufficiency in newborns, according to researchers from the University of Geneva.



© UNIGE

Two brother mice pups from the same litter. The one carrying a mutated Hoxd3 gene is on the left.

High definition pictures

PRESS RELEASE

Geneva | 13th of October 2017

WARNING: embargoed until 16th of October 2017, 19:00 GMT

A family of “architect” genes named Hox coordinates the formation of organs and limbs during embryonic life. Geneticists from the University of Geneva (UNIGE) and the Swiss Federal Institute of Technology in Lausanne (EPFL), Switzerland, have discovered an essential function of one of these genes, Hoxd3, in the development of the gut of suckling mice. Specific mutations in this gene induce defective breast milk assimilation and important, often lethal, growth retardation. In humans, this genetic defect probably contributes to certain types of intestinal insufficiency in premature infants, such as necrotizing enterocolitis of the newborn. The detection of a mutated Hoxd3 gene would enable to identify one of the causes of this disease, which remain unknown to this day. This work was published in the journal *PNAS*.

The Hox genes, which preside over the mammalian construction plan, are involved in the emergence of the digestive system during embryonic development. The group of Denis Duboule, Professor at the Department of Genetics and Evolution of the UNIGE Faculty of Science and at the EPFL, unveiled many functions and modes of action of these architect genes during the last thirty years. “We have observed cases of delayed postnatal growth in pups carrying mutations in the HoxD gene group, and we wanted to determine their precise causes”, says the researcher.

A growth retardation due to a single gene

His team has produced several lines of mice carrying precise and targeted mutations in the HoxD gene cluster. “All of the pups carrying a Hoxd3 gene coding for a dysfunctional protein were severely stunted during the suckling period and displayed an abnormal development of the small intestine”, explains Jozsef Zakany, researcher at the Department of Genetics and Evolution and first author of the study. This gene therefore has a crucial role in the developing gut of suckling mice.

Unlike adults, mammalian babies “eat” milk, incorporating it as undigested droplets. “The Hoxd3 gene appears to promote the maintenance of the intestinal cells responsible for absorbing this nutrient throughout the suckling period. Growth has indeed systematically resumed in surviving mutated animals, reflecting changes in the intestinal wall following weaning”, notes Jozsef Zakany.

From mice to humans

The human and murine genomes are similar, including the genes of the HoxD complex. The same applies to their physiology of nursing. According to the researchers, the presence of a Hoxd3 gene displaying this kind of mutation probably contributes to certain types of intestinal insufficiency in premature infants, such as necrotizing enterocolitis of the newborn. This affliction, which varies from 4 to 22% in babies with a very low birth weight, requires a rapid diagnosis and urgent medical care. "In less than a week, it is possible to test the DNA of infants suspected of having such intestinal insufficiency, to detect a possible mutation of the Hoxd3 gene. This is a path to be explored at the clinical level, as the causes of this disease are still unknown", concludes Denis Duboule.

contact

Jozsef Zakany

+41 22 379 67 88

Jozsef.Zakany@unige.chl

Denis Duboule

+41 22 379 67 71 / +41 21 693 83 38

Denis.Duboule@unige.ch

UNIVERSITÉ DE GENÈVE Communication Department

24 rue du Général-Dufour
CH-1211 Geneva 4

Tel. +41 22 379 77 17

media@unige.ch

www.unige.ch